

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07K 5/062, A61K 38/55</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/13289 (43) International Publication Date: 18 May 1995 (18.05.95)</p>
<p>(21) International Application Number: PCT/GB94/02471 (22) International Filing Date: 10 November 1994 (10.11.94) (30) Priority Data: 9323165.2 10 November 1993 (10.11.93) GB (71) Applicant (for all designated States except US): CHIRO-SCIENCE LIMITED [GB/GB]; Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): MONTANA, John [GB/GB]; Chiroscience Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). DICKENS, Jonathon [GB/GB]; Chiroscience Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). OWEN, David, Alan [GB/GB]; Chiroscience Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). BAXTER, Andrew, Douglas [GB/GB]; Chiroscience Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).</p>		<p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: PEPTIDYL COMPOUNDS AND THEIR THERAPEUTIC USE AS INHIBITORS OF METALLOPROTEINASES</p> <p>(57) Abstract</p> <p>A compound of formula (I), wherein R¹ is C₁₋₆ alkyl, C₂₋₆ alkenyl, aryl, (C₁₋₆ alkyl)aryl or C₁₋₆ alkyl-AR⁹ where A is O, NR⁹ or S(O)_m, where m = 0-2, and the or each R⁹ is H, C₁₋₄ alkyl, aryl or (C₁₋₄ alkyl)aryl; R² is H or C₁₋₆ alkyl; either R⁴ is H or C₁₋₆ alkyl optionally substituted by amino (NH₂), arylamino, protected amino, aryl, di(C₁₋₆ alkyl)amino, mono(C₁₋₆ alkyl)amino, CO₂H, protected carboxyl, carbamoyl, mono(C₁₋₆ alkyl)carbamoyl or di(C₁₋₆ alkyl)carbamoyl, and R⁵ is H or C₁₋₆ alkyl, or NR₄R₅ is pyrrolidino, piperidino or morpholino; R⁶ is optionally-substituted C₃₋₆ cycloalkyl, C₃₋₆ cycloalkenyl, C₁₋₆ alkyl, benzyl, (C₁₋₆ alkoxy)benzyl, benzyloxybenzyl or 3-indolylmethyl; Alk is C₁₋₆ alkyl or C₂₋₆ alkenyl and n = 0 or 1; R⁷ is H or R¹⁰CO where R¹⁰ is C₁₋₄ alkyl, (C₁₋₄ alkyl)aryl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, C₂₋₆ alkenyl or (C₂₋₆ alkenyl)aryl; and R⁸ is H, C₁₋₄ alkyl, (C₁₋₄ alkyl)aryl or aryl; or a salt, solvate or hydrate thereof. Such compounds may be used to treat conditions associated with metalloproteinase, TNF and/or other activity.</p> <div style="text-align: center;"> <p>(I)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PEPTIDYL COMPOUNDS AND THEIR THERAPEUTIC USE
AS INHIBITORS OF METALLOPROTEINASES

Field of the Invention

This invention relates to a novel class of peptidyl derivatives, to processes for their preparation, and to their use in medicine.

5 Background to the Invention

In normal tissues, cellular connective tissue synthesis is offset by extracellular matrix degradation, the two opposing effects existing in dynamic equilibrium. Degradation of the matrix is brought about by the action of proteinases released from resident connective tissue cells and invading inflammatory cells, and is due, in part, to the activity of at least three groups of metalloproteinases. These are the collagenases (interstitial collagenase, MMP-1; PMN collagenase, MMP-8), the gelatinases (gelatinase A, MMP-2, 72kDa-gelatinase, Type IV collagenase; gelatinase B, MMP-9, 92kDa-gelatinase, Type IV collagenase) and the stromelysins (proteoglycanase, MMP-3, stromelysin-1, transin; stromelysin-2, MMP-10; stromelysin-3, MMP-11). Normally, these catabolic enzymes are tightly regulated at the level of their synthesis and secretion and also at the level of their extracellular activity, the latter through the action of specific inhibitors, such as TIMP (tissue inhibitors of metalloproteinase), which form inactive complexes with metalloproteinases, and more general proteinase inhibitors such as α_2 -macroglobulin.

20 The accelerated, uncontrolled breakdown of connective tissues by metalloproteinase-catalysed resorption of the extracellular matrix is a feature of many pathological conditions such as rheumatoid arthritis, osteoarthritis, septic arthritis, corneal, epidermal or gastric ulceration; tumour metastasis or invasion; periodontal disease, proteinuria, coronary thrombosis associated with atherosclerotic plaque rupture and bone disease. Inhibitors may also be useful in preventing the pathological sequelae following a traumatic injury that could lead to a permanent disability. These compounds may also have utility as a means for birth control, by preventing ovulation or implantation. It can be expected that the pathogenesis of such diseases is likely to be modified in a beneficial manner by the administration of metalloproteinase inhibitors, and numerous compounds have been suggested for this purpose [for a general review, see Wahl *et al*, Ann. Rep. Med. Chem. 25: 175-184, Academic Press Inc., San Diego (1990)].

A number of small peptide-like compounds which inhibit metalloproteinases have been described. Perhaps the most notable of these relate to angiotensin-converting enzyme (ACE), where such agents act to block the conversion of the decapeptide angiotensin I to angiotensin II, a potent pressor substance. Compounds of this type are described in EP-A-0012401. Also, related mercaptoamide peptidyl derivatives have shown ACE inhibitor activity *in vitro* and *in vivo* (Weller *et al* (1984) *Biochem. Biophys. Res. Comm.* 125 (1):82-89).

TNF is a cytokine which is produced initially as a cell-associated 28 kD precursor. It is released as an active, 17 kD form (Jue *et al* (1990) *Biochemistry* 29:8371-8377) which can mediate a large number of deleterious effects *in vivo*. When administered to animals or humans it causes inflammation, fever, cardiovascular effects, haemorrhage, coagulation and acute phase responses, similar to those seen during acute infections and shock states. Chronic administration can also cause cachexia and anorexia. Accumulation of excessive TNF can be lethal.

There is considerable evidence from animal model studies that blocking the effects of TNF with specific antibodies can be beneficial in acute infections, shock states, graft versus host reactions and autoimmune disease. TNF is also an autocrine growth factor for some myelomas and lymphomas and can act to inhibit normal haematopoiesis in patients with these tumours.

Preventing the production or action of TNF is, therefore, predicted to be a potent therapeutic strategy for many inflammatory, infectious, immunological or malignant diseases. These include, but are not restricted to, septic shock, haemodynamic shock and sepsis syndrome (Mathison *et al* (1988) *J. Clin. Invest.* 81:1925-1937; Miethke *et al* (1992) *J. Exp. Med.* 175:91-98), post-ischaemic reperfusion injury, malaria (Grau *et al* (1989) *Immunol. Rev.* 112:49-70); mycobacterial infection (Barnes *et al* (1992) *Infect. Imm.* 60:1441-6), meningitis, psoriasis, congestive heart failure, fibrotic disease, cachexia, graft rejection, cancer, autoimmune disease, rheumatoid arthritis, multiple sclerosis, radiation damage, toxicity following administration of immunosuppressive monoclonal antibodies such as OKT3 or CAMPATH-1, and hyperoxic alveolar injury.

Current clinical anti-TNF strategies involve the use of corticosteroids such as dexamethasone, and the use of cyclosporin-A or FK506, which are non-specific

inhibitors of cytokine gene transcription. Phosphodiesterase inhibitors such as pentoxifylline have been shown to be more specific inhibitors of TNF gene transcription (Endres S (1991) Immunol. 72:56-60, Schandene *et al* (1992) Immunol. 76:30-34, Alegre *et al* (1991) Transplantation 52:674-679, Bianco *et al* (1991) Blood 78:1205-1221). Thalidomide has also been shown to inhibit TNF production by leucocytes (Sampajo *et al* (1991) J. Exp. Med. 173:699-703). In experimental settings, anti-TNF monoclonal antibodies, soluble TNF receptors and soluble TNF receptor/immunoadhesins have been shown to specifically inhibit the effects of TNF action (Bagby *et al* (1991) J. Infect. Dis. 163:83-88, Charpentier *et al.* (1991) Presse-med. 20:2009-2011, Silva *et al* (1990) J. Infect. Dis. 162:421-427, Franks *et al* (1991) Infect. Immun. 59:2609-2614, Tracey *et al* (1987) Nature 330:662-664, Fischer *et al* (1992) PNAS USA in press, Lesslauer *et al* (1991) Eur. J. Immunol. 21:2883-2886, Ashkenazi *et al* (1991) PNAS USA 88:10535-10539).

It has recently been shown that the effects of TNF are mediated by two peptides, TNF-a and TNF-b. Although these peptides have only 30% homology with each other, they activate the same receptors and are encoded by immediately adjacent genes. As used herein, the term tumour necrosis factor or TNF therefore means tumour necrosis factor a and peptides having a high degrees of sequence homology with, or substantially similar physiological effects to, TNF-a, for example TNF-b.

Compounds which have the property of inhibiting the action of metalloproteinases involved in connective tissue breakdown such as collagenase, stromelysin and gelatinase have been shown to inhibit the release of TNF both *in vitro* and *in vivo* (Gearing *et al* (1994) Nature 370:555-557, McGeehan *et al* (1994) Nature 370:558-561, WO 93/20047). All of these reported inhibitors contain a hydroxamic acid zinc binding group.

Compounds that inhibit collagenase are disclosed in U.S. Patent No. 4,511,504 issued Apr. 16, 1985, U.S. Patent No. 4,568,666, issued Feb 4, 1986. Compounds of related structure that are claimed to inhibit stromelysin (proteoglycanase) are described in U.S. Patent No. 4,771,037, issued Sept. 13, 1988.

Stromelysin and collagenase inhibitors may have utility in preventing articular cartilage damage associated with septic arthritis. Bacterial infections of the joints can elicit an inflammatory response that may then be perpetuated beyond what is needed for removal of the infective agent, resulting in permanent damage to structural components. Bacterial agents have been used in animal models to elicit an arthritic response with the appearance of proteolytic activities. See Case *et al* (1989) J. Clin. Invest. 84:1731-40 and Williams *et al* (1990) Arth. Rheum. 33: 533-41.

Inhibitors of stromelysin, collagenase and gelatinase may be useful to control tumour metastasis, optionally in combination with current chemotherapy and/or radiation. See Matrisian *et al* (1986) PNAS USA 83:9413-7, Wilhelm *et al* (1987) Ibid. 84:6725-29, Werb *et al* (1989) J. Cell Biol. 109:872-889, Liotta *et al* (1983) Lab. Invest. 49:636-649 and Reich *et al* in Metastasis; Ciba Foundation Symposium, Wiley, Chicester, 1988, pp. 193-210.

Secreted proteinases such as stromelysin, collagenase and gelatinase play an important role in processes involved in the movement of cells during metastatic tumour invasion. Indeed, there is also evidence that the matrix metalloproteinases are over-expressed in certain metastatic tumour cell lines. In this context, the enzyme functions to penetrate underlying basement membranes and allow the tumour cell to escape from the site of primary tumour formation and enter the circulation. After adhering to blood vessel walls, the tumour cells use these same metalloproteinases to pierce underlying basement membranes and penetrate other tissues, thereby leading to tumour metastasis. Inhibition of this process would prevent metastasis and improve the efficacy of current treatments with chemotherapeutics and/or radiation.

These inhibitors should also be useful for controlling periodontal diseases, such as gingivitis. Both collagenase and stromelysin activities have been isolated from fibroblasts derived from inflamed gingiva (Uitto *et al* (1981) J. Periodontal Res. 16:417-424). Enzyme levels have been correlated to the severity of gum disease; Overall *et al* (1987) J. Periodontal Res. 22:81-88.

Proteolytic processes have also been observed in the ulceration of the cornea following alkali burns (Brown *et al* (1969) Arch. Ophthalmol. 81:370-373).

5 Mercapto-containing peptides do inhibit the collagenase isolated from alkali-burned rabbit cornea (Burns (1989) Invest. Ophthalmol 30:1569-1575). Treatment of alkali-burned eyes or eyes exhibiting corneal ulceration as a result of infection with inhibitors of these metalloendoproteinas in combination with sodium citrate or sodium ascorbate and/or antimicrobials may be effective in preventing developing corneal degradation.

10 Stromelysin has been implicated in the degradation of structural components of the glomerular basement membrane (GBM) of the kidney, the major function of which is to restrict passage of plasma proteins into the urine (Baricos *et al* (1989) Biochem. J. 254:609-612). Proteinuria, a result of glomerular disease, is excess protein in the urine caused by increased permeability of the GBM to plasma proteins. The underlying causes of the increased GBM permeability are unknown, but proteinases including stromelysin may play an important role in glomerular diseases. Inhibition of this enzyme may alleviate the proteinura associated with kidney malfunction.

15 Inhibition of stromelysin activity may prevent the rupturing of atherosclerotic plaques, leading to coronary thrombosis. The tearing or rupture of atherosclerotic plaques is the most common event initiating coronary thrombosis. Destabilisation and degradation of the connective tissue matrix surrounding these plaques by proteolytic enzymes or cytokines released by infiltrating inflammatory cells has been proposed as a cause of plaque fissuring. Such tearing of these plaques can cause an acute thrombolytic event, as blood rapidly flows out of the blood vessel. High levels of stromelysin RNA message have been found to be localised to individual cells in atherosclerotic plaques removed from heart transplant patients at the time of surgery

20 (Henney *et al* (1991) PNAS USA 88:8154-8158). Inhibition of stromelysin by these compounds may aid in preventing or delaying the degradation of the connective tissue matrix that stabilises the atherosclerotic plaques, thereby preventing events leading to acute coronary thrombosis.

25 Specific inhibitors of stromelysin and collagenase may be useful as birth control agents. There is evidence that expression of metalloproteinases, including stromelysin and collagenase, is observed in unfertilised eggs and zygotes and at further cleavage stages, and increased at the blastocyst stage of fetal development

30

and with endoderm differentiation (Brenner *et al* (1989) *Genes & Develop.* 3:848-59). By analogy to tumour invasion, a blastocyst may express metalloproteinases in order to penetrate the extracellular matrix of the uterine wall during implantation. Inhibition of stromelysin and collagenase during these early development processes should presumably prevent normal embryonic development and/or implantation in the uterus. Such intervention would constitute a method of birth control. In addition, there is evidence that collagenase is important in ovulation processes. In this example, a covering of collagen over the apical region of the follicle must be penetrated in order for the ovum to escape. Collagenase has been detected during this process and an inhibitor has been shown to be effective in preventing ovulation (Woessner *et al* (1989) *Steroids* 54:491-499). There may also be a role for stromelysin activity during ovulation (Too *et al* (1984) *Endocrin.* 115:1043-1050).

Collagenolytic and stromelysin activities have also been observed in dystrophic epidermolysis bullosa (Kronberger *et al* (1982) *J. Invest. Dermatol.* 79:208-211; Sawamura *et al* (1991) *Biochem. Biophys. Res. Commun.* 184:1003-8). Inhibition of metalloendoproteinases should limit the rapid destruction of connective components of the skin.

In addition to extracellular matrix comprising structural components, stromelysin can degrade other *in vivo* substrates including α_1 -proteinase inhibitor, and may therefore influence the activities of other proteinases such as elastase (Winyard *et al* (1991) *FEBS Letts.* 279(1):91-94). Inhibition of the matrix metalloendoproteinases may potentiate the antiproteinase activity of these endogenous inhibitors.

From recent publications it is evident that several new enzymes of the MMP family have been identified, some of which may be important in disease. Collagenase 3, an enzyme unique to breast carcinoma cells, may have utility in breast cancer (Freije *et al* (1994) *J. Biol. Chem.* 269(24):16766-16773), MT-MMP, another member of the MMP family, has been shown to be a key enzyme in the activation of gelatinase A (Sato *et al* (1994) *Nature* 370:61-65). Gelatinase A is an important enzyme in the growth and metastasis of tumours (such as defined above).

The degradation of b-Amyloid Precursor Protein (APP) has been shown to generate amyloid plaques, a major constituent of the senile plaques, found in patients

with Alzheimers Disease (AD). Two recent publications have identified metalloproteinase enzymes that cleave APP to the amyloid plaque (Abraham *et al* (1994) *Biochemistry* 33:192-199; Huber *et al* (1994) *Biochem. Biophys. Res. Comm.* 201(1):45-53).

5 As will be appreciated by those of skill in the art, the significant proportion of homology between these new enzymes and other MMPs leads to the possibility that a compound that inhibits one enzyme may to some degree inhibit these new enzymes. Therefore, inhibitors encompassed in this invention may be useful in the diseases in which these new enzymes are implicated.

10 An object behind the present invention was to provide compounds which substantially inhibit the release of TNF from cells, and which therefore may be used in the treatment of conditions mediated by TNF. Such uses include, but are not limited to, the treatment of inflammation, fever, cardiovascular effects, haemorrhage, coagulation and acute phase response, cachexia and anorexia, acute infections, shock
15 states, graft versus host reactions and autoimmune disease.

Another object behind this invention was to provide compounds which, possibly in addition to inhibiting TNF release, also inhibit the action of MMPs, and hence may be used in the treatment of patients who suffer from conditions mediated by TNF and/or MMPs.

20 Summary of the Invention

The invention encompasses novel mercaptoalkylpeptidyl compounds of formula (I) which are useful inhibitors of matrix metalloproteinase and/or TNF-mediated diseases including degenerative diseases (such as defined above) and certain cancers.

25 According to one aspect of the invention, novel compounds are of general formula (I), i.e. as shown among the Formulae, below. With respect to formula (I) at least:

R^1 is C_{1-6} alkyl, C_{2-6} alkenyl, aryl, $(C_{1-6}$ alkyl)aryl or C_{1-6} alkyl- AR^9 where A is O, NR^9 or $S(O)_m$, where $m = 0-2$, and the or each R^9 is H, C_{1-4} alkyl, aryl or $(C_{1-4}$ alkyl)aryl;
30

R^2 is H or C_{1-6} alkyl;

X represents NR^4R^5 and

either R^4 is H or C_{1-6} alkyl optionally substituted by amino (NH_2), arylamino, protected amino, aryl, di(C_{1-6} alkyl)amino, mono(C_{1-6} alkyl)amino, CO_2H , protected carboxyl, carbamoyl, mono(C_{1-6} alkyl)carbamoyl or di(C_{1-6} alkyl)carbamoyl, and R^5 is H or C_{1-6} alkyl, or NR_4R_5 is pyrrolidino, piperidino or morpholino;

5 R^3 represents $[Alk]_nR^6$ and R^6 is optionally-substituted C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkyl, benzyl, (C_{1-6} alkoxy)benzyl, benzyloxybenzyl or 3-indolylmethyl, Alk is C_{1-6} alkyl or C_{2-6} alkenyl and $n = 0$ or 1 ;

R^7 is H or $R^{10}CO$ where R^{10} is C_{1-4} alkyl, (C_{1-4} alkyl)aryl, C_{3-6} cycloalkyl, (C_{3-6} cycloalkyl) C_{1-4} alkyl, C_{2-6} alkenyl or (C_{2-6} alkenyl)aryl; and

10 R^8 is H, C_{1-4} alkyl, (C_{1-4} alkyl)aryl or aryl;
or a salt, solvate or hydrate thereof.

Description of the Invention

It will be appreciated that the compounds according to the invention can contain one or more asymmetrically-substituted carbon atoms, for example those marked with an asterisk in formula (I). The presence of one or more of these asymmetric centres in a compound of formula (I) can give rise to stereoisomers, and in each case the invention is to be understood to extend to all such stereoisomers, including enantiomers and diastereomers, and mixtures including racemic mixtures thereof.

20 In the formulae herein, the \sim line is used at a potential asymmetric centre to represent the possibility of *R*- and *S*- configurations, the $<$ line and the line to represent a unique configuration at an asymmetric centre.

As used in this specification, alone or in combination, the term " C_{1-6} alkyl" refers to a straight or branched chain alkyl moiety having from one to six carbon atoms, including, for example, methyl, ethyl, propyl, isopropyl, butyl, t-butyl, pentyl hexyl. Thus, for example, $[Alk]_nR^6$ may be t-butyl.

The term " C_{1-4} alkyl" refers to a straight or branched chain alkyl moiety having from one to four carbon atoms, including, for example, methyl, ethyl, propyl, isopropyl, butyl and t-butyl.

30 The term " C_{2-6} alkenyl" refers to a straight or branched chain alkyl moiety having two to six carbon atoms and having in addition one double bond, of either E

or Z stereochemistry where applicable. This term includes, for example, vinyl, 1-propenyl, 1- and 2-butenyl, and 2-methyl-2-propenyl.

The term "cyclo(C₃₋₆)alkyl" refers to a saturated alicyclic moiety having from three to six carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term "cyclo(C₃₋₆)alkenyl" refers to an alicyclic moiety having from three to six carbon atoms and having in addition one double bond. This term includes, for example, cyclopentenyl and cyclohexenyl.

The term "aryl" means an optionally-substituted phenyl or naphthyl group with the substituent(s) being selected, for example, from halogen, trifluoromethyl, C₁₋₆ alkyl, alkoxy, phenyl and the like. The term "halogen" means fluorine, chlorine, bromine or iodine.

The terms "protected amino" and "protected carboxy" mean amino and carboxy groups which are protected in a manner familiar to those skilled in the art. For example, an amino group can be protected by a benzyloxycarbonyl, *tert*-butoxycarbonyl, acetyl or like groups, or in the form of a phthalimido or like group. A carboxyl group can be protected in the form of a readily-cleavable ester such as the methyl, ethyl, benzyl or *tert*-butyl ester.

The term "alkoxy" refers to a straight chain or branched chain alkoxy group containing a maximum of six carbon atoms, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy or *tert*-butoxy.

Salts of compounds of formula (I) include pharmaceutically acceptable salts, for example acid addition salts derived from inorganic or organic acids, such as hydrochlorides, hydrobromides, *p*-toluenesulphonates, phosphates, sulphates, perchlorates, acetates, trifluoroacetates, propionates, citrates, malonates, succinates, lactates, oxalates, tartrates and benzoates.

Salts may also be formed with bases. Such salts include salts derived from inorganic or organic bases, for example alkali metal salts such as magnesium or calcium salts, and organic amine salts such as morpholine, piperidine, dimethylamine or diethylamine salts.

When the "protected carboxy" group in compounds of the invention is an esterified carboxyl group, it may be a metabolically labile ester of formula CO₂R¹¹

where R¹¹ may be an ethyl, benzyl, phenethyl, phenylpropyl, or α or β -naphthyl, 2,4-dimethylphenyl, 4-tert-butylphenyl, 2,2,2-trifluoroethyl, 1-(benzyloxy)benzyl, 1-(benzyloxy)ethyl, 2-methyl-1-propionyloxypropyl, 2,4,6-trimethylbenzyloxymethyl or pivaloyloxymethyl group.

5 Preferred compounds of the invention include those in which, independently or in any combination:

R¹ represents a C₁₋₄ alkyl or C₁₋₃ alkyl-OH group, for example a 2-methylpropyl (isobutyl) or 2-hydroxyethyl group;

R² represents hydrogen;

10 n is zero;

R⁴ represents hydrogen or a C₁₋₃ alkyl group, for example a methyl group;

R⁵ represents hydrogen or a C₁₋₃ alkyl group, for example a methyl group;

R⁶ represents a benzyl or 3-indolylmethyl group;

R⁷ represents hydrogen or the group R¹⁰CO where R¹⁰ is a C₁₋₄ alkyl group,
15 for example methyl;

R⁸ represents a C₁₋₄ alkyl, aryl or (C₁₋₃ alkyl)aryl group, for example a phenyl or benzyl group;

Particularly preferred compounds according to the invention include:

- N*-[*N*-(Mercaptoacetyl)-L-leucyl]-L-phenylalanine methylamide
- 20 *N*-(Acetylmercaptoacyl)-L-leucyl-L-phenylalanine methylamide
- (*RS*)-2-(Acetylthio)pentanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- (*RS*)-2-(Acetylthio)propanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- (*RS*)-2-(Acetylthio)-3-methylbutanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- (*RS*)-2-(Acetylthio)-2-phenylacetyl-L-leucyl-L-phenylalanine *N*-methylamide
- 25 (*RS*)-2-(Acetylthio)-3-phenylpropanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- (*RS*)-2-(Acetylthio)-4-phenylbutanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- N*-(Acetylmercaptoacyl)-L-threonyl-L-phenylalanine methylamide
- N*-(Acetylmercaptoacyl)-L-leucyl-L-tryptophan methylamide
- (*RS*)-2-Mercaptopentanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- 30 (*RS*)-2-Mercaptopropanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- (*RS*)-2-Mercapto-3-methylbutanoyl-L-leucyl-L-phenylalanine *N*-methylamide
- (*RS*)-2-Mercapto-2-phenylacetyl-L-leucyl-L-phenylalanine *N*-methylamide

(*RS*)-2-Mercapto-3-phenylpropanoyl-L-leucyl-L-phenylalanine *N*-methanamide
(*RS*)-2-Mercapto-4-phenylbutanoyl-L-leucyl-L-phenylalanine *N*-methanamide
N-[*N*-(Mercaptoacetyl)-L-threonyl]-L-phenylalanine methanamide and
N-[*N*-(Mercaptoacetyl)-L-leucyl]-L-tryptophan methanamide

5 Compounds of the general formula (I) may be prepared by any suitable method known in the art and/or by the following processes, which themselves form part of the invention. It will be appreciated that where a particular stereoisomer of formula (I) is required, the synthetic processes described herein may be used with the appropriate homochiral starting material, and/or isomers may be resolved from
10 mixtures using conventional separation techniques (e.g. HPLC).

 The compounds according to the invention may be prepared by the following process. In the description and formulae below, the groups R^1 , R^2 , R^3 , R^6 , R^7 , R^8 , and X are as defined above, except where otherwise indicated. It will be appreciated that functional groups, such as amino, hydroxyl or carboxyl groups, present in the
15 various compounds described below, and which it is desired to retain, may need to be in protected form before any reaction is initiated. In such instances, removal of the protecting group may be the final step in a particular reaction. Suitable protecting groups for such functionality will be apparent to those skilled in the art. For specific details see "Protective Groups in Organic Synthesis", Wiley
20 Interscience, T W Greene, PGM Wuts.

 A process for preparing compounds of general formula (I) comprises deprotecting (for example by hydrolysis) a compound having a general formula [II] which is formula (I) wherein R^7 represents a suitable protecting group (e.g. *tert*-butyl or acetate).

25 It will be appreciated that where a particular stereoisomer of formula (I) is required, this may be obtained by conventional resolution techniques such as high performance liquid chromatography. Where desired, however, appropriate homochiral starting materials may be used in the coupling reaction to yield a particular stereoisomer of formula (I). This is exemplified below.

30 Intermediates of general formula (II) may be prepared by coupling an acid of formula (III) wherein R^7 and R^8 are as defined above, or an active derivative thereof,

with an amine of formula (IV). Active derivatives of acids of formula (III) include, for example, acid anhydrides or acid halides, such as acid chlorides.

The coupling reaction may be performed using standard conditions for amination reactions of this type. Thus, the reaction may be achieved in a solvent, for example an inert organic solvent such as an ether, e.g. a cyclic ether such as tetrahydrofuran, an amide, e.g. a substituted amide such as dimethylformamide, or a halogenated hydrocarbon such as dichloromethane at a low temperature e.g. -30°C to ambient temperature, such as -20°C to 0°C, optionally in the presence of a base, e.g. an organic base such as an amine, e.g. triethylamine or a cyclic amine such as N-methylmorpholine. Where an acid of formula (III) is used, the reaction may additionally be performed in the presence of a condensing agent, for example a diimide such as N,N'-dicyclohexylcarbodiimide, advantageously in the presence of a triazole such as 1-hydroxybenzotriazole. Alternatively, the acid may be reacted with a chloroformate, for example ethyl chloroformate, prior to reaction with the amine of formula (IV).

The amine of general formula (IV) may be prepared by coupling an acid of formula (V), or an active derivative thereof with an amine of formula (VI), followed by removal of any protecting groups.

Active derivatives of acids of formula (V) include, for example, acid anhydrides or acid halides such as acid chlorides as outlined earlier.

Amino acids and their derivatives as depicted by general formulae (V) and (VI) can be obtained in chiral or racemic form. In the chiral form they provide asymmetric building blocks for the chiral synthesis of compounds of general formula (I). Many of these derivatives can be readily obtained from commercially available starting materials using methods known to those skilled in the art. (See "The Practice of Peptide Synthesis" by M. Bodanszky *et al*, Springer Verlag, New York, 1984; WO 92/21360).

Compounds of general formula (II) may be prepared by nucleophilic substitution of compounds of general formula (VII) wherein R¹⁴ represents a suitable leaving group (e.g. a halogen such as bromide, or an alkyl sulphonate ester such as methanesulphonate) with a thiol of general formula (VIII) wherein R⁷ represents a

suitable protecting group (e.g. *tert* butyl or acetate), using standard conditions known to those skilled in the art, as exemplified in WO 90/05719.

Thiols of general formula (VIII) may be obtained from commercially available starting materials using methods known to those skilled in the art. Many
5 thiols of general formula (VIII) are also commercially available.

Compounds of general formula (VII) may be prepared by coupling an acid of general formula (IX) wherein R^{14} and R^8 are as defined above (or suitably protected versions thereof) or an active derivative thereof, with an amine of formula (IV) using similar coupling conditions to those described for the preparation of
10 compounds of formula (II).

Carboxylic acids of the structures depicted in formulae (III) and (IX) can be obtained in chiral or racemic form. Many of these derivatives can be readily obtained from commercially available starting materials, using methods known to those skilled in the art (see WO 90/05719).

15 Compounds of formula (I) may also be prepared by interconversion of other compounds of formula (I). Thus, for example, a compound of formula (I) wherein R^1 is a C_{1-6} alkyl group may be prepared by hydrogenation (using palladium on carbon in suitable solvent, such as an alcohol, e.g. ethanol) of a compound of formula (I) wherein R^1 is a C_{2-6} alkenyl group. A compound of formula (I) wherein
20 R^7 is $R^{10}CO$ may be prepared by acylation (using a suitable acid chloride $R^{10}COCl$, in the presence of a base such as a triethylamine in a suitable solvent, such as a chlorinated solvent, e.g. dichloromethane) of a compound of formula (I) wherein R^7 is H.

Any mixtures of final products or intermediates obtained can be separated on
25 the basis of the physico-chemical differences of the constituents, in known manner, into the pure final products or intermediates, for example by chromatography, distillation, fractional crystallization, or by formation of a salt if appropriate or possible under the circumstances.

The compounds according to the invention exhibit *in vitro* inhibiting activities
30 with respect to stromelysin, collagenase and gelatinase. Compounds according to the invention also exhibit *in vitro* inhibition of TNF release. The activity and selectivity

of the compounds may be determined by use of the appropriate enzyme inhibition test, for example as described in Example A hereinafter.

This invention also relates to a method of treatment for patients (including man and/or mammalian animals raised in the dairy, meat or fur industries or as pets) suffering from disorders or diseases which can be attributed to stromelysin as previously described, and more specifically, a method of treatment involving the administration of the matrix metalloproteinase inhibitors of formula (I) as the active constituents.

Accordingly, the compounds of formula (I) can be used among other things in the treatment of osteoarthritis and rheumatoid arthritis, and in diseases and indications resulting from the over-expression of these matrix metalloproteinases such as found in certain metastatic tumour cell lines.

As mentioned above, compounds of formula (I) are useful in human or veterinary medicine since they are active as inhibitors of TNF and MMPs. Accordingly in another aspect, this invention concerns a method of management (by which is meant treatment or prophylaxis) of disease or conditions mediated by TNF and/or MMPs in mammals, in particular in humans, which method comprises administering to the mammal an effective amount of a compound of formula (I) above, or a pharmaceutically acceptable salt thereof. For this purpose, a compound of formula (I) may be used in the preparation of an agent for the management (by which is meant treatment or prophylaxis) of diseases or conditions mediated by TNF and/or MMPs.

The disease or conditions referred to above include inflammation, fever, cardiovascular effects, haemorrhage, coagulation and acute phase response, cachexia and anorexia, acute infections, shock states, graft versus host reactions and autoimmune disease; and those involving tissue breakdown such as bone resorption, inflammatory diseases, dermatological conditions, tumour growth, angiogenesis and invasion by secondary metastases, in particular rheumatoid arthritis, osteoarthritis, periodontitis, gingivitis, corneal ulceration, tumour growth, angiogenesis and invasion by secondary metastases.

For the treatment of rheumatoid arthritis, osteoarthritis, and in diseases and indications resulting from the over-expression of matrix metalloendoproteinases such

as are found in certain metastatic tumour cell lines or other diseases mediated by the matrix metalloendoproteinases or increased TNF production, the compounds of formula (I) may be administered orally, topically, parenterally, by inhalation, spray or rectally, in dosage unit formulations containing non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs and cats, the compounds of the invention are effective in the treatment of humans.

10 A pharmaceutical composition containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and
15 such compositions may contain one or more agents selected from sweetening agents, flavouring agents, colouring agents and preserving agents, in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be, for example,
20 inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay
25 disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time-delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in US Patents Nos. 4,256,108; 4,166,452; and 4,265,874, to form osmotic therapeutic tablets for control release.

30 Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein

the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long-chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as a polyoxyethylene with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl or n-propyl p-hydroxybenzoate, one or more colouring agents, one or more flavouring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified herein. Sweetening, flavouring and colouring agents may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum

acacia or gum tragacanth, naturally-occurring phosphatides, for example soya bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavouring and colouring agents.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be in a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of formula (I) may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

For topical use, creams, ointments, jellies, solutions or suspensions, etc containing the compounds of formula (I) are employed. For the purposes of this application, topical application includes the use of mouth washes and gargles.

Dosage levels of the order of from about 0.05 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 2.5 mg to about 7 g per patient per day). For example, inflammation may be effectively treated by the administration of from about 0.01 to

50 mg of the compound per kilogram of body weight per day (about 0.5 mg to about 3.5 g per patient per day).

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration of humans may vary from about 5 to about 95 percent of the total composition. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following non-limiting Examples 1-23 illustrate the preparation of compounds of formula (I), using materials including the following Intermediates. Examples A to D illustrate test protocols.

In the Examples, the following abbreviations are used:

DCC Dicyclohexylcarbodiimide

RT Room temperature

EDC 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

THF Tetrahydrofuran

TNF α Tumour necrosis factor α

PMA Phorbol-13-myristate-12-acetate

ELISA Enzyme-linked immunosorbent assay

Intermediate 1 L-Phenylalanine N-methylamide

DCC (3.02 g) was added to a solution of L-phenylalanine-N-benzylcarbamate (4.00 g) and N-hydroxysuccinimide (1.69 g) in dry THF (35 ml) at 0°C. The mixture was stirred at that temperature for 16 h and filtered through Celite. To the filtrate was added aqueous methylamine (40% aq., 5 ml) and the mixture was stirred at RT for 24 h. The reaction mixture was concentrated *in vacuo* and the crude material (6.01 g) was dissolved in ethanol (250 ml). A suspension of 10% palladium on carbon in H₂O (ca 2 ml) was added to the ethanol solution in a bomb. The

mixture was stirred in the presence of hydrogen at 724 kPa (105 psi) for 18 h. The solution was filtered through Celite and the filtrate was evaporated *in vacuo*. The residue was partitioned between 1N hydrochloric acid (110 ml) and dichloromethane (50 ml). The layers were separated and the aqueous layer was extracted with
5 dichloromethane (50 ml). The aqueous layer was basified with 3N sodium hydroxide solution to pH ~ 14 and extracted with dichloromethane (3 x 50 ml). The latter extracts were combined, dried (Na₂SO₄), filtered and evaporated, to give the title compound as a colourless viscous oil which solidified under high vacuum (2.24 g).

TLC R_f 0.15 (5% MeOH-CH₂Cl₂)

10 Similarly prepared was:

Intermediate 2 L-Tryptophan *N*-methanamide

From L-tryptophan-*N*-benzylcarbamate (50 g) as a colourless oil (14.3 g)

TLC R_f 0.30 (5% MeOH-CH₂Cl₂).

Intermediate 3 *N*-[*N*-(Carbobenzyloxy)-L-leucyl]-L-phenylalanine methanamide

15 Intermediate 1 (1.33 g), L-leucine-*N*-benzylcarbamate (1.98 g), *N*-hydroxybenzotriazole (1.21 g) and EDC (1.57 g) were stirred together in dry THF (30 ml) at 0°C for 24 h. Water (10 ml) and dichloromethane (10 ml) was added and the organic solvent was removed *in vacuo*, the aqueous phase was partitioned between dilute hydrochloric acid (1N, 10 ml) and ethyl acetate (20 ml). The organic
20 layer was separated, washed with saturated bicarbonate solution (8%, 20 ml) and brine (20 ml), dried (Na₂SO₄), filtered and evaporated to give an opaque gel. This was redissolved in dichloromethane and evaporated *in vacuo* to give the product as a colourless solid (2.78 g).

TLC R_f 0.60 (EtOAc)

25 Similarly prepared were:

Intermediate 4 *N*-[*N*-(Carbobenzyloxy)-L-leucyl]-L-tryptophan methanamide

From Intermediate 2 (11.4 g) as a white solid (20 g)

TLC R_f 0.46 (5% MeOH-CH₂Cl₂)

Intermediate 5 *N*-[*N*-(Carbobenzyloxy)-L-threonyl]-L-phenylalanine
30 methanamide

From L-threonine-*N*-benzylcarbamate (4.3 g) as a colourless oil (7.7 g)

TLC R_f 0.31 (5% MeOH-CH₂Cl₂)

Intermediate 6 *N*-(*L*-leucyl)-*L*-phenylalanine methylamide

Intermediate 3 (2.78 g) was dissolved in ethanol (140 ml) and then 10% palladium on carbon (0.5 g) (as a slurry in *ca* 3 ml H₂O) was added. The suspension was stirred under a hydrogen atmosphere for 16 h. The mixture was
5 filtered through Celite, washing with dichloromethane (20 ml). Evaporation of the filtrate gave a colourless semi-solid which was dissolved in 1N hydrochloric acid (100 ml) and extracted with dichloromethane (3 x 50 ml), then basified with sodium hydroxide (3N, 50 ml) and extracted with dichloromethane (3 x 50 ml). The latter extracts were combined, dried (Na₂SO₄), filtered and evaporated to give the title
10 compound as a colourless solid (1.66 g).

TLC R_f 0.13 (5% MeOH-CH₂Cl₂)

Intermediate 7 *N*-(*L*-leucyl)-*L*-tryptophan methylamide

A solution of Intermediate 4 (2 g) in cyclohexene (50 ml) and ethanol (50 ml) was treated with 10% palladium on carbon (0.2 g). The mixture was stirred at reflux
15 for 2 h. The solvent was removed *in vacuo* after filtration through Celite and the residue purified by flash column chromatography (SiO₂, 10% MeOH, CH₂Cl₂) to yield a hygroscopic colourless solid (1.05 g).

TLC R_f 0.38 (10% MeOH-CH₂Cl₂)

Similarly prepared was:

Intermediate 8 *N*-(*L*-threonyl)-*L*-phenylalanine methylamide

From Intermediate 5 (6.6 g) as a white solid (3.31 g)

TLC R_f 0.05 (5% MeOH-CH₂Cl₂)

Intermediate 9 Acetylmercaptoacetic acid

Sodium (460 mg) was added to ethanol (20 ml), then thiolacetic acid (1.50
25 ml) was added to the solution. After 5 min, bromoacetic acid (2.78 g) was added and the mixture was stirred at RT for 18 h. The reaction mixture was diluted with dichloromethane (60 ml) and filtered through Celite. The filtrate was evaporated to give a pale yellow oil. The crude product was dissolved in dichloromethane (10 ml), refiltered and evaporated to give the title compound as a pale yellow oil (2.84 g).

30 ¹H NMR (200 MHz; CDCl₃; Ref TMS) 2.41 (s, 3H) and 3.75 (s, 2H)

Intermediate 10 (RS)-2-Bromopentanoic acid

A solution of D,L-*nor*-valine (10 g) and potassium bromide (35.5 g) in aqueous sulphuric acid (1.25M, 130 ml) was treated portionwise at 0°C with solid sodium nitrite (8.8 g) over a period of 1 h. The solution was then allowed to warm to RT and stirred for 2 h. The mixture was extracted with dichloromethane (2 x 100 ml), the combined extracts dried (MgSO₄) and evaporated *in vacuo* to provide the title compound (11.53 g) as a colourless oil.

TLC R_f 0.60 (EtOAc)

Similarly prepared were:

Intermediate 11 (RS)-2-Bromo-3-methylbutanoic acid

From D,L-Valine (10 g) as a colourless oil (11.48 g)

TLC R_f 0.54 (EtOAc)

Intermediate 12 (RS)-2-Bromo-2-phenylacetic acid

From D,L-phenylglycine (10 g) as a pale yellow oil (7.24 g)

TLC R_f 0.58 (EtOAc)

Intermediate 13 (RS)-2-Bromo-3-phenylpropanoic acid

From D,L-phenylalanine (5 g) as a pale yellow oil (6.72 g)

TLC R_f 0.58 (EtOAc)

Intermediate 14 (RS)-2-Bromo-4-phenylbutanoic acid

From D,L-*homo*-phenylalanine (970 mg) as a white solid (1.0 g)

TLC R_f 0.13 (EtOAc)

Intermediate 15 (RS)-2-(Acetylthio)pentanoic acid

A solution of Intermediate 10 (3.0 g) in ethanol (50 ml) was treated with potassium thiolacetate (2.1 g) and the mixture stirred at RT overnight. The solution was evaporated *in vacuo* and the residue partitioned between water (30 ml) and dichloromethane (30 ml). The organic layer was washed with brine (30 ml), dried (MgSO₄) and evaporated *in vacuo* to provide the title compound (2.23 g) as a yellow oil.

TLC R_f 0.55 (EtOAc)

Similarly prepared were:

Intermediate 16 (RS)-2-(Acetylthio)propanoic acid

From (RS)-2-bromopropanoic acid (1.63 g) as a yellow oil (1.38 g)

TLC R_f 0.4 (5% EtOH-CHCl₃)

Intermediate 17 (RS)-2-(Acetylthio)-3-methylbutanoic acid

5 From Intermediate 11 (5 g) as a pale yellow oil (4.1 g)

TLC R_f 0.5 (EtOAc)

Intermediate 18 (RS)-2-(Acetylthio)-2-phenylacetic acid

From Intermediate 12 (3.0 g) as yellow oil (1.64 g)

TLC R_f 0.5 (EtOAc)

10 **Intermediate 19** (RS)-2-(Acetylthio)-3-phenylpropanoic acid

From Intermediate 13 (2.0 g) as a yellow oil (1.41 g)

TLC R_f 0.80 (EtOAc)

Intermediate 20 (RS)-2-(Acetylthio)-4-phenylbutanoic acid

From Intermediate 14 (860 mg) as a tan oil (340 mg)

15 TLC R_f 0.35 (EtOAc)

Example 1 *N*-[*N*-(Acetylmercaptoacetyl)-*L*-leucyl]-*L*-phenylalanine
N-methanamide

Intermediate 6 (113 mg) and Intermediate 9 (52 mg) were dissolved in dry THF (10 ml), *N*-hydroxybenzotriazole (63 mg) was added followed by EDC (82 mg) and the solution was stirred at 0°C for 16 h. The reaction mixture was concentrated *in vacuo* and the residue was partitioned between 1N hydrochloric acid (20 ml) and ethyl acetate (20 ml). The layers were separated and the organic phase was extracted with saturated sodium bicarbonate solution (2 x 20 ml) and brine (20 ml), dried (MgSO₄) and evaporated to give the crude product as a colourless solid (146 mg). Purification by flash column chromatography (8 x 1.5 cm, EtOAc as eluant, loading with CH₂Cl₂) gave the title compound as a colourless solid (110 mg).

TLC R_f 0.35 (EtOAc)

¹H NMR (200 MHz; CDCl₃; Ref TMS): 0.90 (m, 6H), 1.3-1.8 (m, 3H), 2.4 (s, 3H), 2.7 (d, 3H), 2.9-3.3 (m, 2H), 3.5 (s, 2H), 4.2 (m, 1H), 4.6 (m, 1H), 6.1 (m, 1H), 6.45 (br.d, 1H), 6.6 (br.d, 1H), 7.1-7.4 (m, 5H)

Similarly prepared were:

Example 2 (RS)-2-(Acetylthio)pentanoyl-L-leucyl-L-phenylalanine
N-methylamide

From Intermediate 15 (300 mg) and Intermediate 6 (500 mg) as a white solid
(540 mg)

5 TLC R_f 0.63 (5% MeOH-CH₂Cl₂)

C₂₃H₃₅N₃O₄S [449.6]; [MH⁺]=450; [MNa⁺]=472.

Example 3 (RS)-2-(Acetylthio)propanoyl-L-leucyl-L-phenylalanine
N-methylamide

From Intermediate 16 (180 mg) and Intermediate 6 (355 mg) as a white solid

10 (314 mg)

TLC R_f 0.33 (EtOAc)

C₂₁H₃₁N₃O₄S [421.6]; [MH⁺]=422

Example 4 (RS)-2-(Acetylthio)-3-methylbutanoyl-L-leucyl-L-phenylalanine
N-methylamide

15 From Intermediate 17 (330 mg) and Intermediate 6 (553 mg) as a white solid
(570 mg) after recrystallisation from diethyl ether

TLC R_f 0.36, 0.40 (diastereomers) (EtOAc)

C₂₃H₃₅N₃O₄S [449.6]; [MH⁺]=450; [MNa⁺]=47.2

Example 5 (RS)-2-(Acetylthio)-2-phenylacetyl-L-leucyl-L-phenylalanine
N-methylamide

20

From Intermediate 18 (360 mg) and Intermediate 6 (500 mg) as a white solid
(660 mg)

TLC R_f 0.37 (5% MeOH-CH₂Cl₂)

C₂₆H₃₃N₃O₄S [483.6]; [MH⁺]=484, [MNa⁺]=496

25 **Example 6** (RS)-2-(Acetylthio)-3-phenylpropanoyl-L-leucyl-L-phenylalanine
N-methylamide

From Intermediate 19 (420 mg) and Intermediate 6 (553 mg) as a white solid
(410 mg) after recrystallisation from diethyl ether

TLC R_f 0.36, 0.43 (diastereomers) (EtOAc)

30 C₂₇H₃₅N₃O₄S [497.7]; [MH⁺]=498; [MNa⁺]=520

Example 7 (RS)-2-(Acetylthio)-4-phenylbutanoyl-L-leucyl-L-phenylalanine
N-methylamide

From Intermediate 20 (350 mg) and Intermediate 6 (500 mg) as a white solid
(640 mg)

5 TLC R_f 0.39 (5% MeOH-CH₂Cl₂)

C₂₈H₃₇N₃O₄S [511.7]; [MH⁺]=512; [MNa⁺]=534

Example 8 N-[N-(Acetylmercaptoacetyl)-L-threonyl]-L-phenylalanine methylamide

From Intermediate 9 (0.48 g) and Intermediate 8 (1.0 g) as a white solid
(0.83 g)

10 TLC R_f 0.35 (7% MeOH-CH₂-Cl₂)

¹H NMR (200MHz: CDCl₃ + CD₃OD: Ref TMS). 1.1 (d, 3H), 2.4 (s, 3H),
2.7 (d, 3H), 3.1 (m, 2H), 3.6 (d, 2H), 4.2 (m, 2H), 4.6 (m, 1H), 7.2 (m, 5H).

Example 9 N-[N-(Acetylmercaptoacetyl)-L-leucyl]-L-tryptophan methylamide

From Intermediate 9 (220 mg) and Intermediate 7 (500 mg) as a white solid
15 (590 mg)

TLC R_f 0.22 (EtOAc)

C₂₂H₃₀N₄O₄S [446.6]; [MH⁺]=447

Example 10 N-[N-(Mercaptoacetyl)-L-leucyl]-L-phenylalanine methylamide

Example 1 (16 mg) was stirred with saturated ammonium hydroxide solution
20 (200 ml) and methanol (400 ml) at room temperature under nitrogen for 2 h. The
solution was acidified with dilute hydrochloric acid (2N) and the mixture was
extracted with dichloromethane (3 x 40 ml). The combined organic fractions were
dried (Na₂SO₄), filtered and evaporated to give the crude product as a colourless
solid (14.2 mg). Purification by flash column chromatography (1.5 x 10 cm; EtOAc
25 as eluant; loading with CH₂Cl₂) gave the title compound as a colourless solid (6.0
mg).

TLC R_f 0.23 (EtOAc).

¹H NMR (200 MHz; CDCl₃; Ref TMS): 0.90 (d, 3H), 0.93 (d, 3H), 1.48 -
1.71 (m, 3H), 1.86 (t, 1H), 2.74 (d, 3H) 3.02 - 3.17 (m, 4H), 4.40 (m, 1H) 4.62
30 (q, 1H), 6.03 (br m, 1H), 6.82 (br d, 1H), 6.99 (br d, 1H), and 7.12 - 7.37 (m,
5H).

Similarly prepared were:

Example 11 (RS)-3-Mercaptopentanoyl-L-leucyl-L-phenylalanine *N*-methanamide

From Example 2 (200 mg) as a white solid (180 mg)

TLC R_f 0.37 (5% MeOH-CH₂Cl₂)

C₂₁H₃₃N₃O₃S [407.6]; [MH⁺]=408; [MNa⁺]=430

5 **Example 12** (RS)-2-Mercapto-3-methylbutanoyl-L-leucyl-L-phenylalanine
N-methanamide

From Example 4 (200 mg) as a white solid (190 mg)

TLC R_f 0.41 (5% MeOH-CH₂Cl₂)

C₂₁H₃₃N₃O₃S [407.6]; [MH⁺]=408, [MNa⁺]=430

10 **Example 13** (RS)-2-Mercapto-2-phenylacetyl-L-leucyl-L-phenylalanine
N-methanamide

From Example 5 (200 mg) as a white solid (173 mg)

TLC R_f 0.28 [5% MeOH-CH₂Cl₂]

C₂₄H₃₁N₃O₃S [441.6]; [MH⁺]=442, [MNa⁺]=464

15 **Example 14** (RS)-2-Mercapto-3-phenylpropanoyl-L-leucyl-L-phenylalanine
N-methanamide

From Example 6 (202 mg) as a white solid (185 mg)

TLC R_f 0.28 (5% MeOH-CH₂Cl₂)

C₂₅H₃₃N₃O₃S [455.6]; [MH⁺]=456

20 **Example 15** (RS)-2-Mercapto-4-phenylbutanoyl-L-leucyl-L-phenylalanine
N-methanamide

From Example 7 (150 mg) as a white solid (138mg)

TLC R_f 0.20 (5% MeOH-CH₂Cl₂)

C₂₆H₃₅N₃O₃S [469.6]; [MH⁺]=470, [MNa⁺]=492

25 **Example 16** *N*-[*N*-(Mercaptoacetyl)-L-leucyl]-L-tryptophan methanamide

From Example 9 (120 mg) as a white solid (100 mg)

TLC R_f 0.12 (5% MeOH-CH₂Cl₂)

C₂₀H₂₈N₃O₃S [390.5]; [MH⁺]=391

Example 17 *N*-[*N*-(Mercaptoacetyl)-L-threonyl]-L-phenylalanine methanamide

30 From Example 8 (0.3 g) as a white solid (0.09 g)

TLC R_f 0.09 (5% MeOH-CH₂Cl₂)

¹H NMR (200MHz.DMSO) 1.0 (d, 3H), 2.5 (d, 3H), 2.8 (m, 3H), 3.0 (d.d, 1H), 3.2 (m, 2H), 3.9 (m, 1H), 4.2 (d.d, 1H), 4.4 (m, 1H), 5.0 (d, 1H), 7.2 (m, 5H), 7.8 (m, 1H), 8.0 (m, 2H)

Example 18 (RS)-2-Mercaptopropanoyl-L-leucyl-L-phenylalanine *N*-methylamide

- 5 A solution of sodium methoxide (10 mg) in methanol (1.5 ml) was treated with Example 3 (56 mg, 0.133 mmol) and the mixture stirred at room temperature for 1 h. The solvent was evaporated *in vacuo* and the residue partitioned between dichloromethane (15 ml) and 2M hydrochloric acid (10 ml). The aqueous portion was re-extracted with dichloromethane (10 ml) and the combined extracts dried
- 10 (Na₂SO₄) and evaporated *in vacuo* to provide a white solid. Purification by column chromatography (SiO₂, 3% methanol, dichloromethane) provided the title compound (13 mg) as a white solid.

C₁₉H₂₉N₃O₃S [379.5]; MS [MH⁺]=380

TLC R_f 0.32 (5% MeOH-CH₂Cl₂)

15 **Examples 19-23**

Five further compounds that may be made by procedures analogous to those described above are those of formula (I) wherein R² = R⁴ = R⁸ = H, R⁵ = CH₃, R⁶ = ---benzyl, R⁷ = CH₃CO, n = O and:

- R¹ = ◀CH₂SCH₃ (Ex. 19);
- 20 R¹ = ◀CH₂SOCH₃ (Ex. 20);
- R¹ = ◀CH₂SO₂CH₃ (Ex. 21);
- R¹ = ◀CH₂OH (Ex. 22);
- R¹ = ◀CH₂CH₂OH (Ex. 23).

Example A Collagenase inhibition activity

- 25 The potency of compounds of general formula (I) to act as inhibitors of collagenase was determined by the procedure of Cawston and Barrett, (Anal. Biochem., 99:340-345, 1979) whereby a 1 mM solution of the inhibitor being tested or dilutions thereof was incubated at 37°C for 16 hours with collagen and collagenase (buffered with 50 mM Tris, pH 7.6, containing 5 mM CaCl₂, 0.05% Brij 35, 60
- 30 mM NaCl and 0.02% NaN₃). The collagen was acetylated ³H or ¹⁴C-collagen prepared by the method of Cawston and Murphy (Methods in Enzymology 80:711, 1981). The choice of radiolabel did not alter the ability of collagenase to degrade the

collagen substrate. The samples were centrifuged to sediment undigested collagen and an aliquot of the radioactive supernatant removed for assay on a scintillation counter as a measure of hydrolysis. The collagenase activity in the presence of 1mM inhibitor, or a dilution thereof, was compared to activity in a control devoid of inhibitor and the results reported as that inhibitor concentration effecting 50% inhibition of the collagenase (IC_{50}).

Example B Stromelysin inhibition activity

The potency of compounds of general formula (I) to act as inhibitors of stromelysin was determined using the procedure of Cawston *et al* (Biochem. J. 195:159-165, 1981), whereby a 1 mM solution of the inhibitor being tested or dilutions thereof was incubated at 37°C for 16 hours with stromelysin and ^{14}C -acetylated casein (buffered with 50 mM Tris, pH 7.6, containing 5 mM $CaCl_2$, 0.05% Brij 35, and 0.02% NaN_3). The casein was ^{14}C acetylated according to the method described in Cawston *et al*, above. The stromelysin activity in the presence of 1 mM, or a dilution thereof, was compared to activity in a control devoid of inhibitor and the results reported as that inhibitor concentration effecting 50% inhibition of the stromelysin (IC_{50}).








Example C Gelatinase inhibition activity

The potency of the compounds of general formula (I) to act as inhibitors of gelatinase was determined using the procedure of Harris & Krane (Biochem Biophys. Acta 258:566 - 576, 1972), whereby a 1 mM solution of the inhibitor being tested or dilutions thereof was incubated at 37°C for 16 hours with gelatinase and heat-denatured 3H or ^{14}C -acetylated collagen (buffered with 50 mM Tris, pH 7.6, containing 5 mM $CaCl_2$, 0.05% Brij 35 and 0.02% NaN_3). The 3H or ^{14}C gelatin was prepared by denaturing 3H or ^{14}C -collagen produced according to the method of Cawston and Murphy, above, by incubation at 60°C for 30 minutes. Undigested gelatin was precipitated by addition of trichloroacetic acid and centrifugation. The gelatinase activity in the presence of 1 mM, or dilution thereof, was compared to the activity in a control devoid of inhibitor and results reported as that inhibitor concentration effecting 50% inhibition of the gelatinase (IC_{50}).

Example D Inhibition of TNFa production


5 The potency of the compounds of general formula (I) to act as inhibitors of
the production of TNFa was determined using the following procedure. A 1mM
solution of the inhibitor being tested or dilutions thereof was incubated at 37° C in
an atmosphere of 5% CO₂ with U937 cells (human histiosytic lymphoma) suspended
10 in RPM1 1640 medium at a cell density of 2 x 10⁶/ml and stimulated with 50nM
final concentration of PMA. After 18 hours, the supernatant is assayed for the levels
of TNFa using a commercially-available ELISA kit (R & D Systems). The activity
in the presence of 1mM inhibitor or dilutions thereof was compared to activity in a
10 control devoid of inhibitor and results reported as that inhibitor concentration
effecting 50% inhibition of the production of TNFa.

Compounds of the invention have activity, measured by any or all of
Examples A to D.



CLAIMS

1. A compound of formula I



10

wherein R¹ is C₁₋₆ alkyl, C₂₋₆ alkenyl, aryl, (C₁₋₆ alkyl)aryl or C₁₋₆ alkyl-AR⁹ where A is O, NR⁹ or S(O)_m, where m = 0-2, and the or each R⁹ is H, C₁₋₄ alkyl, aryl or (C₁₋₄ alkyl)aryl;

R² is H or C₁₋₆ alkyl;

15

either R⁴ is H or C₁₋₆ alkyl optionally substituted by amino (NH₂), arylamino, protected amino, aryl, di(C₁₋₆ alkyl)amino, mono(C₁₋₆ alkyl)amino, CO₂H, protected carboxyl, carbamoyl, mono(C₁₋₆ alkyl)carbamoyl or di(C₁₋₆ alkyl)carbamoyl, and R⁵ is H or C₁₋₆ alkyl, or NR₄R₅ is pyrrolidino, piperidino or morpholino;

20

R⁶ is optionally-substituted C₃₋₆ cycloalkyl, C₃₋₆ cycloalkenyl, C₁₋₆ alkyl, benzyl, (C₁₋₆ alkoxy)benzyl, benzyloxybenzyl or 3-indolylmethyl;

Alk is C₁₋₆ alkyl or C₂₋₆ alkenyl and n = 0 or 1;

R⁷ is H or R¹⁰CO where R¹⁰ is C₁₋₄ alkyl, (C₁₋₄ alkyl)aryl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, C₂₋₆ alkenyl or (C₂₋₆ alkenyl)aryl; and

R⁸ is H, C₁₋₄ alkyl, (C₁₋₄ alkyl)aryl or aryl;

25

or a salt, solvate or hydrate thereof.

2. A compound of claim 1, wherein R¹ is alkyl, alkenyl, aryl or alkylaryl; and R⁷ and R⁸ are each H.

3. A compound of claim 1, selected from

N-[N-(mercaptoacetyl)-L-leucyl]-L-phenylalanine methylamide

30

N-(acetomercaptoacetyl)-L-leucyl]-L-phenylalanine methylamide

(RS)-2-(acetylthio)pentanoyl-L-leucyl-L-phenylalanine N-methylamide

(RS)-2-(acetylthio)propanoyl-L-leucyl-L-phenylalanine N-methylamide

- (RS)-2-(acetylthio)-3-methylbutanoyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-(acetylthio)-2-phenylacetyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-(acetylthio)-3-phenylpropanoyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-(acetylthio)-4-phenylbutanoyl-L-leucyl-L-phenylalanine N-methylamide
 5 N-(acetylmercaptoacyl)-L-threonyl-L-phenylalanine methylamide
 N-(acetylmercaptoacyl)-L-leucyl-L-tryptophan methylamide
 (RS)-2-mercaptopentanoyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-mercaptopropanoyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-mercaptop-3-methylbutanoyl-L-leucyl-L-phenylalanine N-methylamide
 10 (RS)-2-mercaptop-2-phenylacetyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-mercaptop-3-phenylpropanoyl-L-leucyl-L-phenylalanine N-methylamide
 (RS)-2-mercaptop-4-phenylbutanoyl-L-leucyl-L-phenylalanine N-methylamide
 N-[N-(mercaptoacetyl)-L-threonyl]-L-phenylalanine methylamide and
 N-[N-(mercaptoacetyl)-L-leucyl]-L-tryptophan methylamide.
 15 4. A pharmaceutical composition for use in therapy, comprising a compound of
 any of claims 1 to 3 and a physiologically-acceptable diluent or carrier.
 5. Use of a compound of any of claims 1 to 3, for the manufacture of a
 medicament for the prevention or treatment of a condition associated with matrix
 metalloproteinase or TNF activity.
 20 6. Use according to claim 5, wherein the condition is selected from
 inflammation, fever, cardiovascular effects, haemorrhage, coagulation and acute
 phase response, cachexia and anorexia, acute infections, shock states, graft versus
 host reactions, autoimmune disease, malaria, reperfusion injury, meningitis,
 psoriasis, rheumatoid arthritis, multiple sclerosis and tumours.

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/GB 94/02471

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K5/062 A61K38/55

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ALMQUIST, RONALD G. ET AL 'Development of peptidomimetic inhibitors of a newly isolated atrial peptide-degrading enzyme' 1992, PEPT.: CHEM. BIOL., PROC. AM. PEPT. SYMP., 12TH (1992), MEETING DATE 1991, PAGES 791-2. EDITOR(S): SMITH, JOHN A.; RIVIER, JEAN E. PUBLISHER: ESCOM, LEIDEN, NETH. CODEN: 57XGA9 see example 9; table 1 ---	1
X	J. LABELLED COMPD. RADIOPHARM. (1993), 33(11), 1065-78 CODEN: JLCRD4; ISSN: 0362-4803, 1993 BORMANS, G. ET AL 'Synthesis and labeling characteristics of technetium-99m-mercaptoacetyltriptides' see the whole document ---	1
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

9 March 1995

Date of mailing of the international search report

16. 03 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Fuhr, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 94/02471

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,88 06890 (RES CORP TECHNOLOGIES INC) 22 September 1988 see claims; examples ---	1,2,4-6
P,X	WO,A,94 07481 (MERCK & CO INC ;HAGMANN WILLIAM K (US); KOPKA IHOR E (US)) 14 April 1994 see claims; examples ---	1-6
A	EP,A,0 273 689 (BEECHAM GROUP PLC) 6 July 1988 see claims; examples -----	1-6

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 94/02471

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-8806890	22-09-88	AT-T- 106087	15-06-94
		DE-D- 3889726	30-06-94
		DE-T- 3889726	13-10-94
		EP-A- 0358660	21-03-90
		JP-T- 2502537	16-08-90

WO-A-9407481	14-04-94	AU-B- 5292193	26-04-94

EP-A-0273689	06-07-88	AU-B- 599736	26-07-90
		AU-A- 8292687	01-09-88
		DE-A- 3774858	09-01-92
		IE-B- 60751	10-08-94
		JP-A- 63215660	08-09-88
		US-A- 4937243	26-06-90
		US-A- 5109000	28-04-92
